

# Immunization Requirements for School-Age Children

3 Years	4 Years	5 Years	11-15 Years	16 Years and older
<a href="#">DTaP</a> – 4 doses	<a href="#">*DTaP</a> – 4 or 5 doses	<a href="#">*DTaP</a> – 4 or 5 doses	<a href="#">*DTaP</a> – 4 or 5 doses	<a href="#">*DTaP</a> – 4 or 5 doses
<a href="#">Polio (IPV)</a> – 3 doses	<a href="#">*Polio (IPV)</a> – 3 or 4 doses	<a href="#">*Polio (IPV)</a> – 3 or 4 doses	<a href="#">*Polio (IPV)</a> – 3 or 4 doses	<a href="#">*Polio (IPV)</a> – 3 or 4 doses
<a href="#">MMR</a> – 1 dose	<a href="#">MMR</a> – 2 doses	<a href="#">MMR</a> – 2 doses	<a href="#">MMR</a> – 2 doses	<a href="#">MMR</a> – 2 doses
<a href="#">*Hib</a> – 3 or 4 doses	<a href="#">*Hib</a> – 3 or 4 doses	<a href="#">*Hib</a> – no requirement	<a href="#">Tdap</a> – 1 dose	<a href="#">Tdap</a> – 1 dose
<a href="#">Hepatitis A</a> - 2 doses	<a href="#">Hepatitis A</a> - 2 doses	<a href="#">Hepatitis A</a> - 2 doses	<a href="#">Hepatitis A</a> -2 doses	<a href="#">Hepatitis A</a> 2 doses
<a href="#">Hepatitis B</a> – 3 doses	<a href="#">Hepatitis B</a> – 3 doses	<a href="#">Hepatitis B</a> – 3 doses	<a href="#">Hepatitis B</a> – 3 doses	<a href="#">Hepatitis B</a> – 3 doses
<a href="#">*Varicella</a> – 1 dose	<a href="#">*Varicella</a> – 2 doses	<a href="#">*Varicella</a> – 2 doses	<a href="#">*Varicella</a> – 2 doses	<a href="#">*Varicella</a> – 2 doses
<a href="#">*Pneumococcal (PCV)</a> – 4 doses	<a href="#">*Pneumococcal (PCV)</a> – 4 doses	<a href="#">*Pneumococcal (PCV)</a> – no requirement	<a href="#">Meningococcal ACWY (MenACWY)</a> – 1 dose (at 11-12 years of age)	<a href="#">*Meningococcal ACWY (MenACWY)</a> – 2 doses for 16 yrs and older

\*DTaP- Fifth dose not needed if dose #4 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Polio - Fourth dose not needed if dose #3 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Hib - last dose must be on or after 12 months of age-not required after age 5 years; 3 or 4 doses depending on brand used

\*HepB - Alternative schedule for 11-15 year olds – 2 doses of adult Hep B

\*PCV - last dose must be on or after 12 months of age-not required after age 5 years

\*Varicella - no doses needed if history of **diagnosed** Chickenpox disease

\*Meningococcal ACWY – if initial dose given at 16 yrs or older, will only need one dose

# Immunization Requirements for Childcare/Preschool Children

By Age 3 months	By Age 5 months	By Age 7 months	By Age 12 months	By Age 16 months	By Age 19 months	By Age 4 years	By Age 5 years
<a href="#">DTaP</a> 1 dose	<a href="#">DTaP</a> 2 doses	<a href="#">DTaP</a> 3 doses	<a href="#">DTaP</a> 3 doses	<a href="#">DTaP</a> 4 doses	<a href="#">DTaP</a> 4 doses	<a href="#">*DTaP</a> 4 or 5 doses	<a href="#">*DTaP</a> 4 or 5 doses
<a href="#">Polio (IPV)</a> 1 dose	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 2 doses	<a href="#">Polio (IPV)</a> 3 doses	<a href="#">*Polio (IPV)</a> 3 or 4 doses	<a href="#">*Polio (IPV)</a> 3 or 4 doses
<a href="#">Hib</a> 1 dose	<a href="#">Hib</a> 2 dose	<a href="#">Hib</a> 2 doses	<a href="#">*Hib</a> 3 doses	<a href="#">*Hib</a> 3 or 4 doses	<a href="#">*Hib</a> 3 or 4 doses	<a href="#">*Hib</a> 3 or 4 doses	<a href="#">Hib</a> – no requirement
<a href="#">Hepatitis B</a> 1 dose	<a href="#">Hepatitis B</a> 2 doses	<a href="#">Hepatitis B</a> 2 doses	<a href="#">Hepatitis B</a> 2 doses	<a href="#">Hepatitis B</a> 2 doses	<a href="#">Hepatitis B</a> 3 doses	<a href="#">Hepatitis B</a> 3 doses	<a href="#">Hepatitis B</a> 3 doses
<a href="#">Pneumococcal (PCV)</a> 1 dose	<a href="#">Pneumococcal (PCV)</a> 2 doses	<a href="#">*Pneumococcal (PCV)</a> 3 doses	<a href="#">*Pneumococcal (PCV)</a> 4 doses	<a href="#">*Pneumococcal (PCV)</a> 4 doses	<a href="#">*Pneumococcal (PCV)</a> 4 doses	<a href="#">*Pneumococcal (PCV)</a> 4 doses	<a href="#">*Pneumococcal (PCV)</a> – no requirement
			<a href="#">Hepatitis A</a> 1 dose	<a href="#">Hepatitis A</a> 1 dose	<a href="#">Hepatitis A</a> 2 doses	<a href="#">Hepatitis A</a> 2 doses	<a href="#">Hepatitis A</a> 2 doses
				<a href="#">MMR</a> 1 dose	<a href="#">MMR</a> 1 dose	<a href="#">MMR</a> 2 doses	<a href="#">MMR</a> 2 doses
				<a href="#">Varicella</a> 1 dose	<a href="#">Varicella</a> 1 dose	<a href="#">Varicella</a> 2 doses	<a href="#">Varicella</a> 2 doses

\*DTaP-Fifth dose not needed if dose #4 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Polio-Fourth dose not needed if dose #3 given after 4<sup>th</sup> B-day and 6 months from last dose

\*Hib- May have fewer doses if start series late; last dose must be on or after 12 months of age; not required after age 5 years; 3 or 4 doses depending on brand used

\*PCV- May have fewer doses if start series late; last dose must be on or after 12 months of age; not required after age 5 years

\*Varicella-no doses needed if history of **diagnosed** Chickenpox disease